arXiv:1110.3165v1 [cond-mat.mes-hall] 14 Oct 2011

Two-phase stretching of molecular chains

Alexander V. Savin,[1] Mikhail A. Mazo,[1] Irina P. Kikot,[1] and Alexey V. Onufriev[2]
[1] *Semenov Institute of Chemical Physics, Russian Academy of Sciences, Moscow 119991, Russia*
[2] *Departments of Computer Science and Physics,*
2160C Torgersen Hall, Virginia Tech, Blacksburg, VA 24061, USA

While stretching of most polymer chains leads to rather featureless force-extension diagrams, some, notably DNA, exhibit non-trivial behavior with a distinct plateau region. Here we propose a unified theory that connects force-extension characteristics of the polymer chain with the convexity properties of the energy vs. stretch profile of its individual monomer subunits. Namely, if the inter-atomic interaction potentials within the monomer are such that its is described by a uniformly convex (concave down) energy function, the stretching of the entire chain is single phase and uniform. However, if the monomer stretching energy as a function of extension has a non-convex (concave up) region, the stretching of the polymer chain separates into two phases: weakly and strongly stretched monomers. We use simplified planar and 3D polymer models to illustrate the basic principles of the proposed model. Specifically, we show that when the secondary structure of a polymer is mostly due to weak non-covalent interactions, the stretching is two-phase, and the force-stretching diagram has the characteristic plateau. This scenario is realized in α-helix, consistent with earlier observations based on all-atom molecular dynamics simulations. Using a coarse-grained model of the DNA with realistic potentials, we show explicitly that the double helix exhibits the two-phase behaviour under stretching, in agreement with single molecule experiments. The predicted end points of the force-extension plateau are in close quantitative agreemnt with experiment, while a reasonable qualitative agreement is obtained for the tension value at the plateau. Detailed analysis demonstrtates that breaking of Watson-Crick bonds is not necessary for the existence of the plateau, although some of the bonds do break as the double-helix extends.

I. INTRODUCTION

Early on, the mechanical properties of DNA attracted the interest of both physicists and biologists, because of their importance to numerous biological processes, such as DNA transcription, gene expression and regulation, and DNA replication. Besides, mechanical properties are essential while constructing nanodevices containing DNA.

In the past decade new methods appeared which made it possible to stretch single molecule and measure a stretching force with high precision. In these experiments one end of the molecule is attached to the surface, another one is attached to a latex bead which is pulled by optical or magnetic tweezers [1–4]. The main result of such experiments is a force-extension diagram. Complex behavior has been revealed by elasticity studies of individual double-stranded DNA molecules. Under normal conditions several different regimes have been observed [5]. Until distance between molecule ends is less than 0.8 of its contour length the force (chain tension) increases linearly with extension increasing, according to Hooke's law. It should be mentioned that rigidity in this regime is rather small, $2 \div 3$ pN force is sufficient to stretch a molecule for 90% of its contour length [1]. The force in this regime has purely entropic nature. When the end-to-end distance is close to the contour length the force increases significantly [1, 2]. This effect is explained by intrinsic elasticity, i.e. related with structure deformation, which begins to contribute significantly when the contour length of the molecule is approached. Here elastic response is due to both entropic factor and intrinsic elasticity. So when end-to-end distance is less than contour length molecule behaviour is described by ordinary persistent model [5]. When the molecule is subjected to forces of $65 \div 120$ pN, it suddenly changes form, stretching up to 70% beyond its canonical B-form contour length. Various models for the structure of this so-called S-form DNA have been proposed, but its nature remains obscured. When higher forces are applied molecule divides into two strands and afterwards covalent bonds break, chemical structure is disrupted.

Single molecule stretching experiments have been carried out also for polypeptide molecules. [6–8]. The common polypeptide has complex secondary structure, and its force-extension diagram is rather complex; but similar diagrams with typical force-extension plateau have been observed experimentally for synthetical alpha-helices [7] and myosin molecules [6].

Force-extension plateau observed in single DNA molecule experiments is usually explained by either gradual unzipping of the double helix (force-induced tension) [3, 4, 9, 10], or cooperative transition of the whole structure into a new form called S-form where hydrogen bonds between complementary bases remain unbroken. The latter is supported by the fact that transition occurs in a narrow range of forces (about several pN) and is reversible [1, 2], as well as by existing of the second transition under significantly higher forces (about $150 \div 300$ pN), which is attributed to non-equilibrium melting [11]. The version of the force-induced tension is supported firstly by estimates of free energy which reveal that under room temperature the melted state is more favorable due to entropic effects [9], secondly by coincidence of force-extension diagrams for double-stranded and single-stranded DNA on certain range [9], thirdly by experiments which visualize double- and single-stranded segments using specific fluorescent labels [12]. Computer simulations also don't enable to determine the nature of the force-extension plateau: different investigation support existing of S-form DNA, where base pairing remains intact [1, 13, 14], as well as possibility of force-induced melting [15, 16].

In the case of polypeptide force-extension plateau is attributed to the alpha-helix unwinding [6, 17, 18]. each site is supposed to have two stable states which have different free energy — winded and unwinded. Then this representation can be rather easily described by Izing type models [19, 20]. This phenomenon is called helix-coil transition.

Assumption that two phase (strongly and slightly stretched) occur in polymer molecule stretching process has been suggested previously for DNA molecule [1, 19–21]. Usually the strongly stretched state was supposed to consist of melted base pairs, where hydrogen bonds are ruptures [9, 19, 20]. But a model has been supposed [21] where hydrogen bonds remain in strongly stretched state, which is stable and corresponds to an energy minimum.

The paper considers mechanical stretching of three different molecular chains: plane zigzag chain, helical chain and DNA double helix in coarse-grained representation. Our results reveal that for linear polymers non-uniform stretching can take place only when non-valent interactions contribute significantly to its secondary structure stabilization. It is the case for polypeptide α-helix which is stabilized by hydrogen bonds and for DNA double helix which is stabilized by stacking interactions and hydrogen bonds between complementary nitrogen bases. Thus plateau in a force-extension diagram is a consequence of relation between non-valent and valent interactions.

It should be noted that current paper consider only mechanical energy (instead of free energy which is considered in most papers). However even consideration of this constituent only enables to explain a typical force-extension diagram. Besides, computer simulation of DNA double helix dynamics under the temperature 300 K, performed within the framework of coarse-grained representation, indicates that main result remains correct independently of thermal fluctuations. In the stretched state hydrogen bonds are weakened significantly, so certain hydrogen bonds may rupture due to thermal fluctuations. However these hydrogen bonds ruptures can not be considered as a cause for the plateau on the force-extension diagram.

Assumption about two-phase stretching have been proposed previously [6, 17, 21], but the stretched state was always considered as stable, that is corresponding to the free energy minimum. The current paper demonstrates that such an assumption is not necessary for two-phase scenario of stretching (the right end of the plateau may be not the minimum of the energy function).



Figure 1: Dependence of uniformly stretched molecular chain energy E (curve 1) from relative longitudinal step extension $\Delta\rho$. The straight line (line 2), connecting points $\Delta\rho_1$ and $\Delta\rho_2$ and being the tangent in these points, describes a convex hull of the function $E(\Delta\rho)$, $\Delta\rho_0$ stands for the inflection point.

II. THE GENERAL MECHANISM

Aforementioned features of a force-extension diagram for DNA molecule and polypeptide alpha-helix can be explained in a generalized way. The chain structure is stabilized by interactions of two types: valent and non-valent. The former describe bond, angle and torsion deformations. The latter describe interactions between atoms which are not connected by rigid valent interactions. These interactions include electrostatic and van der Waals interactions, hydrogen bonds in polypeptide alpha-helix and stacking interactions between neighboring base pairs in DNA. The main feature of the non-valent interactions potentials $V(r)$ is existing of the inflection point which is a left end of an interval where deformation energy increasing slows significantly. If non-valent interactions contribute significantly to uniformly stretched chain energy $E(\Delta\rho)$ ($\Delta\rho \geq 0$ is relative extension), then the function $E(\Delta\rho)$ also may have an inflection point $\Delta\rho_0$. In this case the function $E(\Delta\rho)$ is no longer convex downwards; its convex hull can be obtained by connecting two points $(\Delta\rho_1, E(\Delta\rho_1))$ and $(\Delta\rho_2, E(\Delta\rho_2)))$ $(0 < \Delta\rho_1 < \Delta\rho_0 < \Delta\rho_2)$ by a common tangent – see Fig. 1. This line describes more favorable way of stretching when one part of the chain is in strongly stretched state with extension $\Delta\rho_1$, and another one is in slightly extended state with the extension $\Delta\rho_2$. While this non-uniform stretching chain extension is due to increasing of the strongly stretched sites, so the tension remains constant. Such non-uniform stretching scenario explains completely the existing of the force-extension plateau.



Figure 2: Scheme of the zigzag molecular chain.

III. STRETCHING OF THE 2-D ZIGZAG MOLECULAR CHAIN

Let's explain the mechanism of the non-homogeneous two-phase stretching on example of 2D zigzag chain depicted on the Fig. 2. Let a molecular chain have a form of plane zigzag with fixed x-step h_x and y-step h_y. All atoms of the chain are situated in the xy-plane. In the equilibrium state n-th atom has coordinates $x_n = nh_x$, $y_n = [1 + (-1)^n]h_y/2$. A zigzag is described by the distance between neighboring sites $\rho_0 = (h_x^2 + h_y^2)^{1/2}$ and zigzag angle θ_0 [x-step is $h_x = \rho_0 \sin(\theta_0/2)$, y-step is $h_y = \rho_0 \cos(\theta_0/2)$].

The 2D zigzag energy is given by the following hamiltonian:

$$H = \sum_n \{\frac{1}{2}M(\dot{x}_n^2 + \dot{y}_n^2) + V(\rho_n) + U(\theta_n) + W(r_n)\}, \quad (1)$$

with M being an atom mass, (x_n, y_n) − n-th atom coordinates. The potential $V(\rho_n)$ describes the energy of interaction between neighboring n-th and $(n+1)$-th atoms depending from the distance $\rho_n = (a_n^2 + b_n^2)^{1/2}$, where $a_n = x_{n+1} - x_n$, $b_n = y_{n+1} - y_n$. This potential corresponds to the deformation energy of a valent bond between neighboring atoms. The potential $U(\theta_n)$ describes the deformation energy of the angle between three atoms $(n-1)$-th, n-th and $(n+1)$-th, which is given by

$$\theta_n = \arccos[-(a_n a_{n-1} + b_n b_{n-1})/\rho_{n-1}\rho_n].$$

The last term $W(r_n)$ describes weak non-valent interaction between n-th and $(n+2)$-th atoms (second neighbors), with $r_n = [(x_{n+2} - x_n)^2 + (y_{n+2} - y_n)^2]^{1/2}$ being a distance between these atoms.

Let's consider dimensionless zigzag model where equilibrium bond length is $\rho_0 = 1$, zigzag angle is $\theta_0 = \arccos(-1/3) = 109.47^\circ$ (so that x-step is $h_x = 0.8165$, y-step is $h_y = 0.5774$).

Rigid bond deformation is given by

$$V(\rho) = \frac{1}{2}K(\rho - \rho_0)^2, \quad (2)$$

with bond rigidity being $K = 2$.

Angle deformation is given by

$$U(\theta) = \epsilon_1(\cos\theta - \cos\theta_0)^2, \quad (3)$$

where $\epsilon_1 \geq 0$ is angle deformation coefficient.

The energy of non-valent interactions between second neighbors is given by the potential

$$W(r) = \epsilon_0 \left[\left(\frac{r_0 - d}{r - d} \right)^6 - 1 \right]^2, \quad (4)$$

where $\epsilon_0 \geq 0$ is interaction energy, $r_0 = 2h_x = 1.633$ is equilibrium length, $d = 0.5$ is a diameter of the inner hard core. For the sake of simplicity equilibrium length r_0 was supposed to correspond to the equilibrium value of the angle θ_0, although it may be not the case. The potential $W(r)$ has a form which is typical for non-valent interactions, it may be used for description of hydrogen bonds and van der Waals interactions.

A similar model has been used in the papers [22–25] for analysis of the longitudinal soliton waves motion. It should be mentioned that many polymer macromolecules have 2D zigzag form. For example, in the three-dimensional space a polyethylene (PE) molecule has a stable plane conformation of *trans-zigzag*. The 2D zigzag form is also usual for hydrogen bond chains $\cdots$X–H$\cdots$X–H$\cdots$X–H$\cdots$ of halides, where X=F, Cl, Br, I.

Let's consider zigzag deformation during its homogeneous stretching. If we denote the extension of the zigzag x-step by $\Delta x > 0$, then for finding stretched zigzag form one has to solve the minimization problem over ρ

$$E(\rho) = V(\rho) + U(\theta) + W(r) \to \min : \rho \quad (5)$$

with fixed value of x-step $h_x + \Delta x$ [a distance between second neighbors is $r = 2(h_x + \Delta x)$, value of the angle θ is defined by bond length ρ]. Solving the problem (5) yields the dependence of zigzag energy (per 1 site) from the relative x-step extension Δx.

Let's fix a rigidity of the second neighbors interactions $\kappa = 1$ (interaction energy is $\epsilon_0 = \kappa(r_0 - d)^2/72 = 0.0178$), while angle deformation energy coefficient ϵ_1 will be changed.

With $\epsilon_1 = 0$ the secondary structure of the molecular chain is defined only by non-valent interactions between second neighbors. Dependence of the angle θ and energy E from the relative uniform extension Δx are shown on the Fig. 3. Zigzag stretching is accompanied by monotonous angle increasing. When $\Delta x = 0.1836$ the zigzag straightens out completely in a line (zigzag angle is $\theta = 180^\circ$).

One can see from the Fig. 3 (b) that the function $E(\Delta x)$ is not convex downward. But if one connects two points $(\Delta x_1, E(\Delta x_1))$ and $(\Delta x_2, E(\Delta x_2))$ ($\Delta x_1 = 0.0370$, $\Delta x_2 = 0.2056$) on the curve $(\Delta x, E(\Delta x))$ by the straight line which is a tangent of the curve in these two points, one gets a convex function. This connecting line describes more energetically favorable way of the zigzag stretching when one part of the zigzag is in slightly extended state with the x-step $h_x + \Delta x_1$, and another part is in strongly expanded state with the x-step $h_x + \Delta x_2$, when the zigzag has straightened out in a line.

Dependence of the stretching force $F = dE/d\Delta x$ from the relative extension Δx is shown on the Fig. 3 (c).



Figure 3: Dependence of (a) angle θ, (b) energy E and (c) force $F = dE/d\Delta x$ in the zigzag from its relative uniform extension Δx. Angle deformation energy coefficient is $\epsilon_1 = 0$. The straight line 1 describe a convex hull of the function $E(\Delta x)$.

One can see from the figure that if non-uniform way of stretching is taken into account the force-extension dependence $F(\Delta x)$ has a typical form with constant central part. When $0 \le \Delta x \le \Delta x_1$ and $\Delta x_2 \le \Delta x$, that is during uniform stretching of the zigzag, tension increases monotonously. When $\Delta x_1 < \Delta x < \Delta x_2$, that is during non-uniform stretching, tension remains constant. A typical plateau appears, which has been discovered earlier in experiments dedicated to stretching of alpha-helix [18] and DNA double helix [1, 2].

The deformation energy coefficient ϵ_1 being increased, the typical form of dependencies $E(\Delta x)$ and $F(\Delta x)$ is preserved up to maximal value $\epsilon_1 = 0.015$. Here non-uniform stretching takes place when $0.0710 < \Delta x < 0.1925$. Energy benefit of the non-uniform stretching becomes negligible in this case.

When $\epsilon_1 > 0.015$ the function $E(\Delta x)$ becomes convex downwards — see the Fig. 4. Here only uniform stretching of the zigzag can take place. When $\epsilon_1 = 0.02$ zigzag angle increases monotonously and under $\Delta x = 0.2343$ reaches the maximal value $\theta = 180°$ (zigzag straightens out in a line). The function $E(\Delta x)$ is convex downwards, so the tension F increases monotonously with zigzag extension.



Figure 4: The dependence of (a) angle θ, (b) energy E and tension $F = dE/d\Delta x$ of the zigzag from a relative extension Δx. Angle deformation energy coefficient is $\epsilon_1 = 0.02$.

For verifying obtained results let's consider the changing of zigzag structure consisting from $N = 400$ atoms while stretching. For this purpose for each value of relative extension $\Delta x \ge 0$ one has to solve the minimization problem

$$\sum_{n=1}^{N-1} V(\rho_n) + \sum_{n=2}^{N-1} U(\theta_n) + \sum_{n=1}^{N-2} V(r_n) \to \min_{\{x_n,y_n\}_{n=1}^N} \qquad (6)$$

with a condition of fixed ends:

$$x_1 \equiv -\frac{1}{2}(N-1)\Delta x, \quad x_N = (N-1)h_x + \frac{1}{2}(N-1)\Delta x.$$

If $\{x_n, y_n\}_{n=1}^N$ is a solution of the minimization problem (6), then distribution of the longitudinal extension in the chain is given by the function $\Delta x_n = x_{n+1} - x_n - h_x$, and distribution of y-step is given by the function $h_n = |y_{n+1} - y_n|$. Dependence of distribution Δx_n and h_n from the value of local extension Δx are shown on the Fig. 5 and 6.

One can see from the Fig. 5 that when $\epsilon_1 = 0$ a distribution of x-step extension Δx_n and y-step h_n along



Figure 5: The dependence of distribution of (a) longitudinal extension Δx_n and (b) y-step h_n in the stretched zigzag from $N = 400$ atoms with fixed ends from the extension per site Δx. Angle deformation energy coefficient is $\epsilon_1 = 0$.

the chain corresponds to the behaviour of the function $E(\Delta x)$. When $0 \le \Delta x \le 0.037$ and $\Delta x \ge 0.2056$ the uniform stretching takes place, while when $0.037 < \Delta x < 0.2056$ the non-uniform one is observed. Here end parts of the zigzag are in strongly expanded state with relative extension $\Delta x_n = 0.2056$, while the central part is in slightly stretched state with relative extension $\Delta x_n = 0.037$. On the domain boundary gradual transition from one state to another is observed. Extension increasing, central slightly stretched part monotonously decreases and vanishes with $\Delta x = 0.2056$. Further zizgzag is stretched uniformly.

Similar scenario is observed when the limit value $\epsilon_1 = 0.015$ is considered. Here uniform stretching takes place when $0 \le \Delta x \le 0.0710$ and $\Delta x \ge 0.1925$. With $0.0710 < \Delta x < 0.1925$ the zigzag has two-phase structure — end parts are in strongly expanded state and central part is in slightly extended state.

When $\epsilon_2 = 0.02$, see Fig. 6, the zigzag stretching is always uniform.

So when angle rigidity is zero or rather small stretching of a zigzag chain always has the two-phase scenario. As a consequence, tension remains constant in some range of extensions. Such scenario is due to the typical form of the interaction between second neighbors (4), which have an inflection point. So two-phase stretching can be expected to occur in molecular chain whose secondary structure is formed by weak non-valent interactions. From the other hand, if the secondary structure is due to angle and torsion deformation, then stretching will be uniform, that is all sites will be stretched equally. Such a scenario is typical for polyethylene (PE) trans-zigzag and polytetrafluoroethylene (PTFE) helix.



Figure 6: The dependence of distribution of (a) longitudinal extension Δx_n and (b) y-step h_n in the stretched zigzag from $N = 400$ atoms with fixed ends from the extension per site Δx. Angle deformation energy coefficient is $\epsilon_1 = 0.02$.

IV. STRETCHING OF THE ALPHA-HELIX

Let's consider helical molecular chain with atoms having equilibrium coordinates

$$\mathbf{R}_n^0 = (R_0 \cos(n\phi_0), R_0 \sin(n\phi_0), nh_z), \qquad (7)$$

with $n = 0, \pm 1, \pm 2, ...$ being atom number, R_0 – helical radius, ϕ_0 and h_z – angle and longitudinal helix period.

Chain energy is described by the hamiltonian

$$H = \sum_n \{\frac{1}{2} M(\dot{\mathbf{R}}_n, \dot{\mathbf{R}}_n) + V(\rho_n) + U(\theta_n) + Z(\delta_n) + W(r_n)\}, \qquad (8)$$

where 3D vector $\mathbf{R}_n(t)$ stands for the position of the n-th site (peptide group) of the helix in the moment t, M is site mass. The potential $V(\rho_n)$ gives an energy of interaction between neighbor sites n-th and $(n+1)$-th, depending from the distance $\rho_n = |\mathbf{u}_n|$, where $\mathbf{u}_n = \mathbf{R}_{n+1} - \mathbf{R}_n$. The deformation energy of the angles is described by $U(\theta_n)$, where θ_n is an angle with a vertex in the n-th site, $\cos(\theta_n) = -(\mathbf{u}_n, \mathbf{u}_{n-1})/\rho_{n-1}\rho_n$. The third potential $Z(\delta_n)$ describes deformation energy of the torsion around n-th bond. The torsion cosine is given by $\cos(\delta_n) = (\mathbf{v}_n, \mathbf{v}_{n+1})/|\mathbf{v}_n||\mathbf{v}_{n+1}|$, where the vector$\mathbf{v}_n = \mathbf{u}_{n-1} \times \mathbf{u}_n$. The potential $W(r_n)$ stands for the energy of the n-th hydrogen bond connecting n-th and $(n+3)$-th sites, its length is given by $r_n = |\mathbf{R}_{n+3} - \mathbf{R}_n|$.

Again for the sake of simplicity we consider dimensionless model of helix, where equilibrium bond length is $\rho_0 = 1$, angular step is $\phi_0 = 100°$, equilibrium angle is $\theta_0 = \arccos(-1/3) = 109.47°$. Then [26] dimensionless helix radius is

$$R_0 = \rho_0 \cos(\theta_0/2)/[1 - \cos(\phi_0)] = 0.4919,$$

z-step is

$$\Delta z = \rho_0 \sqrt{|\cos(\phi_0) + \cos(\theta_0)|/[1 - \cos(\phi_0)]} = 0.6572,$$

equilibrium torsion is $\delta_0 = \arccos(0.2395)$, equilibrium hydrogen bond length is $r_0 = 2.0322$.

Bond deformation energy is described by the potential (2) with the rigidity $K = 10$. Angle deformation energy is described by the potential (3) with the deformation energy coefficient $\epsilon_1 = 1$. Torsion deformation energy is given by the potential

$$W(\delta) = \epsilon_2[\cos(\delta) - \cos(\delta_0)]^2, \tag{9}$$

where $\epsilon_2 \geq 0$ – is deformation energy coefficient. Hydrogen bond energy is given by the potential (4) with a parameter $d = 0.7$ (dimensionless group length) and energy $\epsilon_0 = \kappa(r_0 - d)^2/72 = 0.0246$, where rigidity $\kappa = 1$.

Scheme of the alpha-helix is shown on the Fig. 7 (a). Similar helix models have been used in the papers [26, 27] for analysis of ultrasonic soliton motion.



Figure 7: (a) Scheme of the alpha-helix. Helix sites are shown on their equilibrium positions, solid lines denote rigid valent bonds while dotted lines are for soft hydrogen bonds. (b) A zone of the transition between strongly and slightly stretched parts of the helix.

Let's consider changing of the helix structure while stretching. If z-step increases by $\Delta z > 0$, then to define helical structure changing one has to solve a problem of finding conditional minimum by two parameters: radius R and angular step ϕ

$$E(R, \phi, h_z + \Delta z) = V(\rho) + U(\theta) + W(r) + Z(\delta) \to \min_{R,\phi} \tag{10}$$

with the fixed value of z-step $h_z + \Delta z$. Solving the minimization problem (10) yields the dependence of helix energy E (per one step) from a value of z-step relative extension Δz. Firstly the case $\epsilon_2 = 0$ is considered. Without torsional rigidity helix form is stabilized only by hydrogen bonds. Dependencies of the energy E, angular step ϕ and radius R of the helix from its relative longitudinal extension Δz are shown on the Fig. 8. The helix being stretched its angular step monotonously increases and reaches the maximal value $\phi = 180°$ when $\Delta z = 0.1647$ (the helix is unwinded into plane zigzag).

One can see from the Fig. 8 (a) that the function $E(\Delta z)$ is not convex downward. But if one connects two points $(\Delta z_1, E(\Delta z_1))$ and $(\Delta z_2, E(\Delta z_2))$ ($\Delta z_1 = 0.0643$, $\Delta z_2 = 0.1694$) on the curve $(\Delta x, E(\Delta x))$ by the straight line which is a tangent of the curve in these two points, it gets a convex function. This connecting line describes more energetically favorable way of the zigzag stretching where one part of the zigzag is in slightly extended state with the z-step $h_z + \Delta z_1$, and another part is in strongly expanded state with the z-step $h_z + \Delta z_2$, when the helix is unwinded into the plane zigzag.

When $\Delta z_1 < \Delta z < \Delta z_2$ the tension of the helix $F = dE/d\Delta z$ remains constant. Here the dependence $F(\Delta z)$ has typical view with constant central part, as well as for the zigzag.

If torsion deformation energy coefficient ϵ_2 is increased this typical view of the dependence $E(\Delta z)$ is preserved until the maximum value $\epsilon_2 = 0.0015$ is reached. When $\epsilon_2 > 0.0015$ the function $E(\Delta z)$ becomes convex downwards – see fig. 9. Here only uniform stretching of the helix will take place. When $\epsilon_2 = 0.002$ helix stretching is accompanied by monotonous increasing of its angular step ϕ, which reaches its maximal value $\phi = 180°$ when $\Delta z = 0.1727$ (the helix has a form of plane zigzag).

To verify our results about two-phase stretching let's



Figure 8: Dependence of (a) the energy E, (b) angular step ϕ and (c) radius R of the helix from its relative longitudinal extension Δz. Torsion deformation energy coefficient is $\epsilon_2 = 0$. The line 1 shows a convex hull of the function $E(\Delta z)$.

consider changing of the structure of stretched helix, consisting from $N = 400$ sites, under tension. For each value of relative extension $\Delta z \geq 0$ one has to solve the minimization problem

$$\sum_{n=1}^{N-1} V(\rho_n) + \sum_{n=2}^{N-1} U(\theta_n) + \sum_{n=2}^{N-2} Z(\delta_n) + \sum_{n=1}^{N-3} W(r_n) \rightarrow \min : \{(R_n, \phi_n, z_n)\}_{n=1}^{N} \quad (11)$$

with the condition of fixed ends

$$z_1 \equiv -\frac{1}{2}(N-1)\Delta z, \quad z_N \equiv (N-1)h_z + \frac{1}{2}(N-1)\Delta z.$$

The vector (R_n, ϕ_n, z_n) gives cylindric coordinates of n-th site.

If $\{(R_n, \phi_n, z_n)\}_{n=1}^{N}$ is a solution of minimization problem (11), then distribution of longitudinal extension along the helix is given by the function $\Delta z_n = z_{n+1} - z_n - h_z$. Dependence of distribution Δz_n from relative extension Δz is shown on the Fig. 10.



Figure 9: Dependence of (a) the energy E, (b) angular step ϕ and (c) radius R of the helix from its relative longitudinal extension Δz. Torsion deformation energy coefficient is $\epsilon_2 = 0.002$.

It can be seen from the Fig. 10, that if $\epsilon_2 = 0$ the distribution of longitudinal extension Δz_n along the chain completely corresponds to behaviour of the function $E(\Delta z)$. If $\Delta z \leq 0.065$ the chain is stretched uniformly, while if $0.07 \leq \Delta z \leq 0.165$ non-uniform stretching is observed. End parts of the chain are in strongly stretched state while the central part is in slightly stretched state. On the ends the chain has a form of plane zigzag, in the central part it has a form of 3D helix. A bound of transition between strongly stretched central part to slightly end part is shown on the Fig. 7 (b). Extension increasing, central part is monotonously decreased and vanishes when $\Delta z = 0.170$. If extension is larger the chain is stretched uniformly. When $\epsilon_2 = 0.0015$ non-uniform stretching is observed when $0.105 \leq \Delta z \leq 0.155$. If $\epsilon_2 = 0.002$ only uniform stretching can be observed.

Thus independently of torsional rigidity stretching of the helix always has two-phase scenario. As a consequence, tension remains constant in some range of extensions. Such scenario is due to the typical form of the hydrogen bond potential (4), which have an inflection point.



Figure 10: Dependence of distribution of longitudinal extension Δz_n in a stretched helix of $N = 400$ sites from relative extension Δz with (a) $\epsilon_2 = 0$, (b) $\epsilon_2 = 0.0015$ and (c) $\epsilon_2 = 0.002$.

The investigation reveals that extension of polypeptide alpha-helix will be non-uniform, what is confirmed by all-atomic computer simulations [18]. It should be noted that polymer helix which have no hydrogen bonds, such as PTFE helix, will be stretched uniformly, tension increases when extension increases.

V. STRETCHING OF THE DNA DOUBLE HELIX

For simulating DNA double helix stretching a more complex coarse-grained model is used. Each nucleotide consists of six grains – three for sugar-phosphate backbone and three for nucleobase. The model is described in details in the paper [28]. A scheme of the model is depicted on the Fig. 11. In this case a hamiltonian of the system has following form:

$$H = E_k + E_v + E_b + E_a + E_t + E'_{hb} + E_{st} + E_{el} + E_{vdW}. \quad (12)$$



Figure 11: Grains involved into valent interactions. Blue lines denote valent (harmonic) bonds, red arcs mark valent angles. The circles marked as N stand for the atoms N9 on A, G bases and N9 on T, C bases (no grain is situated on these atoms, their coordinates are calculated additionally from base grains positions).

The first term is a kinetic energy of the system.

$$E_k = \sum_{n=1}^{N} \sum_{i=1}^{12} \frac{1}{2} M_i \dot{\mathbf{r}}_{n,i}^2,$$

where i is a number of a grain in n-th base pair.

Terms E_v, E_a, E_t describe energy of correspondingly valent bonds, angles and torsions deformation. These potentials have rather common form: bond deformation energy is calculated by the formula

$$U_{\alpha\beta}(r) = \frac{1}{2} K_{\alpha\beta}(r - R_{\alpha\beta})^2,$$

rigidity coefficients $K_{\alpha\beta}$ and equilibrium values $R_{\alpha\beta}$ are different for different grains. Angle deformation energy has a form

$$U_a(\theta) = \epsilon_a(\cos\theta - \cos\theta_a)^2,$$

where values of coefficient ϵ_a and equilibrium angle differ for different angle types. Torsion deformation energy is described by the potential

$$U_t = \epsilon_t(1 - \cos(\phi - \phi_0)),$$

where values of coefficient ϵ_t and equilibrium torsion angle ϕ_0 differ for different torsions.

Rotational axis of torsions are shown on the Fig. 11. The third term E_b in the hamiltonian (12) describes deformation energy of a nitrogen base. The nitrogen base is rather rigid chemical structure what is modeled by rigid harmonic potentials which keep four points – grain C1 and three points on a nucleobase – near one plane.

The terms E'_{hb} and E_{st} describes interactions between nitrogen bases (including stacking and hydrogen bonds). Since a nitrogen base is rather rigid structure, we can calculate coordinates of all initial atoms from positions of three grains. Thus all-atomic AMBER [29] Coulombic and van der Waals potentials are used to mimic hydrogen bonds and stacking.

The last two terms E_{el}, E_{vdW} describe electrostatic and van der Waals interactions between backbone grains. Solvent is treated implicitly, using the Generalized Born approach [30, 31].

Solvent molecules make hydrogen bonds rupture much easier (after rupture nitrogen bases form hydrogen bonds with water molecules). To take into account this effect one has to weaken hydrogen bonds between complementary bases. For this purposes modified potential $E'_{hb} = c_0 E_{hb}$ is used, where E_{hb} is potential of interactions between complementary bases from all-atomic AMBER potential system [29], and $c_0 = 0.4$ is normalizing factor. Values of parameters and details of potential calculation can be found in the paper [28].

For definiteness homogeneous sequence polyA-polyT is considered. Such double helix can be considered as quasi-one-dimensional crystal with the elementary cell being one nucleotide pair of the double helix. In the ground state each successive nucleotide pair is obtained from its predecessor by translation along the z-axis by step Δz and by rotation around the same axis through helical step $\Delta\phi$. Thus, the energy of the ground state is a function of 38 variables: $\{\mathbf{x}_{1,j}\}_{j=1}^{12}$ and $\Delta\phi$, Δz, where $\mathbf{x}_{1,j} = (x_{1,j,1}, x_{1,j,2}, x_{1,j,3})$ is a radius-vector of j-th grain of the first nucleotide pair.

To find the ground state of the homogeneous double helix under tension, it is necessary to solve minimization problem over 37 variables

$$E = E_v + E_b + E_a + E_t + E'_{hb} + E_{st} + E_{el} + E_{vdW} \rightarrow \min : \{\mathbf{x_j}\}_{j=1}^{12},\ \Delta\phi, \quad (13)$$

under the fixed value of longitudinal step Δz (for calculating the energy one should keep in mind that summation is taken over only one base pair and neighboring base pairs are obtained from it by rotation and translation). As a result, one can obtain the dependence of the homogeneous state energy (per one base pair) E on the longitudinal step Δz – see Fig. 12 (a). One can see from the figure that basic minimum is reached with z-step $\Delta z_0 = 3.352$Å, which corresponds to the ground state of B conformation of double helix DNA.

View of ground state under two typical values of z-step $\Delta z = \Delta z_1$ and $\Delta z = \Delta z_2$ is shown on the Fig. 13. It can be seen from the figure that in strongly stretched state the helix is unwinded more, but all the hydrogen bonds remain – only inclination angle of base pair increases.



Figure 12: Dependence of (a) ground state energy E, (b) tension $F = dE/d\Delta z$ (s) hydrogen bonds energy $-E'_{hb}$ and (d) neighbor basepair stacking interaction energy $-E_{st}$ from longitudinal step Δz of polyA DNA ground state. Lines 1,2,3 give dependencies for uniform stretching while lines 2 and 5 give dependencies for non-uniform stretching of DNA double helix.

One can see from the Fig. 12(a) that when $\Delta z > \Delta z_0$ the function $E(\Delta z)$ is not convex downward. But if one connects two points $(\Delta z_1, E(\Delta z_1))$ and $(\Delta z_2, E(\Delta z_2))$ ($\Delta z_1 = 3.75$Å, $\Delta z_2 = 6.15$Å) on the curve $(\Delta x, E(\Delta x))$ by the straight line which is a tangent of the curve in these two points, he gets a convex function. This connecting line (Fig. 12, line 2) describes more energetically favorable way of the helix stretching where one part of the zigzag is in slightly extended state with the z-step Δz_1, and another part is in strongly expanded state with the z-step Δz_2. The fig. 12 shows that such scenario consumes less energy than uniform stretching of all sites.

Dependence of hydrogen bond energy from extension is shown on the Fig. 12 (c). Under tension hydrogen bonds weaken. When z-step $\Delta z = \Delta z_2$ hydrogen bond weakens approximately twice. One can see from the Fig. 12 (d)



Figure 13: View of ground state of (a) slightly stretched (longitudinal step is $\Delta z = \Delta z_1$) and (b) strongly stretched (z-step is $\Delta z = \Delta z_2$) polyA DNA double helix.



Figure 14: View of ground state of (a) slightly stretched (longitudinal step is $\Delta z = \Delta z_1$) and (b) strongly stretched (z-step is $\Delta z = \Delta z_2$) polyA DNA double helix under the temperature $T = 300$ K.

that weakening of stacking interactions also occurs while stretching. Thus we can conclude that both hydrogen bonds and stacking interactions are responsible for nonconvexity of the function $E(\Delta z)$.

Under the room temperature $T = 300$K weakened hydrogen bonds rupture due to thermal fluctuation – see Fig. 14. Therefore in the case of non-uniform stretching DNA macromolecule consists of two parts: one part is slightly stretched double helix where hydrogen bonds remain [see Fig. 14 (a)], another one is strongly stretched double helix where some hydrogen bonds are ruptured [see Fig. 14 (b)]. Exactly this behaviour is observed in all-atomic DNA stretching simulations[15, 32]. In the papers [3, 4, 9, 10] existing of the plateau in force-extension diagram is attributed to the rupture of hydrogen bonds. But our analysis reveals that it isn't the case. The plateau is observed also in the absence of thermal fluctuations, when hydrogen bonds are only weakened, but not yet ruptured.

Force-extension dependence $F = dE(\Delta z)/d\Delta z$ is shown on the Fig. 12 (b). The force is seen to increase monotonously in the range of z-step from Δz_0 to Δz_1 (curve 3). When the value of z-step Δz_1 is reached uniform stretching became non-uniform one. In the range $\Delta z_1 \le \Delta z \le \Delta z_2$, when non-uniform stretching takes place, tension remains constant (curve 5). When the value Δz_2 is reached stretching again becomes uniform, and tension increases monotonously with Δz increasing (curve 4).

The plateau in force-extension diagram has been observed previously in single molecule stretching experiments [1, 2], the plateau was reported to occur in the range $1.1 < \Delta z/\Delta z_0 < 1.7$. So, having $\Delta z_0 = 3.34$, experiments estimate plateau range as 3.67Å$< \Delta z <$ 5.68Å, which agree well with current results, obtained within the framework of CG representation 3.75Å$< \Delta z <$ 6.15Å [see Fig. 12 (b)]. Value of there transition force 0.2 eV/Å= 320 pN is somewhat higher than experimental estimates, this matter is discussed below. It should be noted that this value should be sequence-depended.

Molecular dynamics simulation have demonstrated that the scenario of non-uniform stretching is preserved under the room temperature 300K, only range of the plateau increases slightly, but the value of force coincide with the value under T=0 K.

In the reported experiments the value of tension in the plateau depends significantly from conditions, such as temperature, salt, sequence and especially from a way of fixing ends. The typical value of force is usually reported to be about 65 ÷ 70 pN [1, 2, 12]; however one should keep in mind that this value was obtained while stretching by the same type ($3'$ or $5'$) ends, while the other two ends were free. On the contrary, the simulations reported here corresponds to uniform stretching by all four ends. Experimentally, larger values of the plateau tension were reported in this case: 105 ÷ 120pN [12, 33], although that is is still less than half of the 320 pN predicted by our model. However, since many DNA properties are strongly sequence-dependent, only semi-quantitative comparison of our homogeneous model with the above experiment (that used a specific, non-uniform sequence) is possible. It is known from other experiments that, for example, pure poly(dG-dC) and poly(dA-dT) DNA sequences, pulled by two of the same type ends, yield tension values at the plateau that differ by a factor of two[33].

It should be mentioned that difference between results of all-atomic simulations and experimental values is of the same order [1, 13]. Therefore numerical agreement with experimental values can be considered as reasonable.

VI. CONCLUSIONS

The investigation demonstrates that existing of non-valent interactions can result in appearing an inflection point in the dependence of homogeneous chain energy from extension. As a consequence a non-uniform way of stretching becomes energetically favorable, when one part of a chain is stretched slightly and another part is stretched strongly. Extension of the whole chain is due to increasing of fraction of strongly stretched sites, so the tension remains constant. Such way of non-uniform stretching explains the plateau in a force-extension diagram.

The non-uniform stretching is observed in stretching simulations of both polypeptide alpha-helix and DNA double helix. Obtained results enable to give a general explanation of the force-extension plateau observed both for DNA double helix [1, 2, 13] and polypeptide alpha-helix [6, 7, 18] in single molecule stretching experiments, as well as in corresponding computer simulations. In the case of alpha-helix gradual unwinding of the helix have been supposed to occur while stretching [6]. However suggested models of this event were based upon the assumption that strongly stretched state is stable, that is corresponds to the free energy minimum. Our investigations reveal that it may be not the case.

In the case of DNA molecule the force-extension plateau is explained either by cooperative transition into the S-form conformation [1, 13, 14] or by force-induced melting [3, 4, 9, 10]. Our results allow to conclude that stretching is accompanied by gradual transition from slightly stretched to strongly stretched state, but the strongly stretched state isn't stable. However some hydrogen bonds ruptures may occur due to thermal fluctuations. Obtained estimates of force-extension plateau are in reasonable agreement with experiments.

[1] P. Cluzel, A.Lebrun, A. Heller, R. Lavery, J.-L. Viovy, D. Chatenay, and F. Caron, Science **271**, 792 (1996).

[2] S. B. Smith, Y. Cui, and C. Bustamante, Science **271**, 795 (1996).

[3] M. C. Williams, K. Pant, I. Rouzina, and R. L. Karpel, Spectroscopy **18**, 203 (2004).

[4] M. J. McCauley and M. C. Williams, Biopolymers **91**, 265 (2008).

[5] C. Bustamante, S. B. Smith, J. Liphardt, and D. Smith, Current Opinion in Structural Biology **10**, 279 (2000).

[6] I. Schwaiger, C. Sattler, D. R. Hostetter, and M. Riff, Nature Letters **1**, 232 (2002).

[7] R. Afrin, I. Takahashi, K. Shiga, and A. Ikai, Biophysical J. **96**, 1105 (2009).

[8] F. Ritort, J. Phys.: Condens. Matter **18**, R531 (2006).

[9] I. Rouzina and V. A. Bloomfield, Biophys. J. **80**, 882 (2001).

[10] I. Rouzina and V. A. Bloomfield, Biophys. J. **80**, 894 (2001).

[11] M. Rief, H. Clausen-Schaumann, and H. E. Gaub, Nat Struct Mol Biol **6**, 346 (1999).

[12] J. van Mameren, P. Gross, G. Farge, P. Hooijman, M. Modesti, M. Falkenberg, G. J. L. Wuite, and E. J. G. Peterman, Proc. Natl. Acad. Sci. USA **106**, 18231 (2009).

[13] A. Lebrun and R. Lavery, Nucleic Acids Research **24**, 2260 (1996).

[14] R. Lavery and A. Lebrun, Genetica **106**, 75 (1999).

[15] H. Li and T. Gisler, Eur. Phys. J. E **30**, 325 (2009).

[16] J.-S. Chen, H. Teng, and A. Nakano, Finite Elements in Analysis and Design **43**, 346 (2007).

[17] M. Rief, J. M. Fernandez, and H. E. Gaub, Phys. Rev. Lett. **81**, 4764 (1998).

[18] F. C. Zegarra, G. N. Peralta, A. M. Coronado, and Y. Q. Gao, Physical Chemistry Chemical Physics **11**, 4019 (2009).

[19] B. H. Zimm and J. K. Bragg, J. Chem. Phys. **31**, 526 (1959).

[20] T. Hill, J. Chem. Phys. **30**, 383 (1959).

[21] P. Cizeau and J.-L. Viovy, Biopolymers **42**, 383 (1997).

[22] A. V. Zolotaryuk, P. L. Christiansen, and A. V. Savin, Phys. Rev. E **54**, 3881 (1996).

[23] L. I. Manevitch and A. V. Savin, Phys. Rev. E **55**, 4713 (1997).

[24] A.V.Savin, L.I.Manevitch, P. L. Christiansen, and A. V. Zolotaryuk, Uspekhi Fizicheskih Nauk **169**, 255 (1999).

[25] A. V. Savin, L. I. Manevich, P. L. Christiansen, and A. V. Zolotaryuk, Physics-Uspekhi. **42**, 245 (1999).

[26] P. L. Christiansen, A. V. Zolotaryuk, and A. V. Savin, Phys. Rev. E **56**, 877 (1997).

[27] A. V. Savin and L. I. Manevitch, Phys. Rev. E **61**, 7065 (2000).

[28] A. V. Savin, M. A. Mazo, I. P. Kikot, L. I. Manevitch, and A. V. Onufriev, Phys. Rev. B **83**, 245406 (2011).

[29] W. D. Cornell, P. Cieplak, C. I. Bayly, I. R. Gould, K. M. Merz, Jr. D. M. Ferguson, D. C. Spellmeyer, T. Fox, J. W. Caldwell, and P. A. Kollman, J. Am. Chem. Soc. **117**, 5179 (1995).

[30] W. C. Still, A. Tempczyk, R. C. Hawley, and T. Hendrickson, J. Am. Chem. Soc. **112**, 6127 (1990).

[31] A. Onufriev, D. A. Case, and D. Bashford, Journal of Computational Chemistry **23**, 1297 (2002).

[32] S.A.Harris, Z.A.Sands, and C.A.Laughton, Biophys. J. **88**, 1684 (2005).

[33] H. Clausen-Schaumann, M. Rief, C. Tolksdorf, and H. E. Gaub, Biophysical Journal **78**, 1997 (2000).